

Medical Record

Tracking Tool

for Newborn Hearing Screening

Name: _____

DOB: _____

ALWAYS ASK ABOUT PARENTAL CONCERNS RE: HEARING

Hospital Hearing Screen

- PASS
- REFER: *Diagnostic Testing*

_____ Date: _____


_____ Notes: _____

Diagnostic Testing Results

- PASS
- HEARING LOSS DIAGNOSED:
 - Birth to Three* Referred to: _____
 - ENT Referral* _____
 - Genetic Testing* _____
 - Ophthalmological evaluation* _____

Birth to Three

- ELIGIBLE: *Review & Sign IFSP*
- NOT ELIGIBLE
 - Track according to rescreening*
 - Monitor developmental issues*



All newborns in Connecticut should be screened before leaving the hospital but in the absence of screening results, these newborns need immediate screening.

Newborns at risk for hearing loss

- NICU admission > 48 hours
- Stigmata or other findings consistent with syndrome known to include hearing loss
- Family history of childhood sensorineural hearing loss
- Craniofacial anomalies
- In-utero infection (for examples see list)

Syndromes associated with progressive hearing loss:

- Neurofibromatosis, Osteopetrosis, Usher's Syndrome

In-utero infections:

- Cytomegalovirus, Rubella, Syphilis, Herpes, or Toxoplasmosis

Tracking Tool

Medical Record

for Newborn Hearing Screening



Age	1 mo	2 mo	4 mo	6 mo	9 mo	12 mo	15 mo	18 mo	24 mo	3 yr
Responds to sound (e.g. starting, blinking)	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Responds to loud noises	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Turns head to voice	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Turns to sound	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Imitates vocalizations	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Understands a few words	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Points to objects when named	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Points to body parts when asked	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Imitates words	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Sings simple songs	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Asks why questions	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Parent/Caregiver concerns and hearing addressed at this visit:

Conditions that require evaluation every 3 to 6 months

- Cranio-facial abnormalities
- Family history
- Head trauma
- In utero infection (for examples see list)
- Mild/unilateral loss previously detected
- Neonatal indicators (for examples see list)
- Neurodegenerative disorder (for examples see list)
- Ototoxic medication (for examples see list)
- Postnatal infection including bacterial meningitis
- Recurrent/Persistent OME
- Speech/language delay
- Stigmata or syndromes associated with hearing loss (for examples see list)

In-utero Infections:

- Cytomegalovirus, Rubella, Syphilis, Herpes, or Toxoplasmosis

Neonatal Indicators:

- Hyperbilirubinemia requiring transfusion, pulmonary hypertension associated with mechanical ventilation, use of ECMO

Neurodegenerative Disorders:

- Hunter syndrome, sensory motor neuropathies, such as Friedreich's ataxia and Charcot-Marie-Tooth syndrome

Ototoxic Medications:

- Chemotherapeutic agents or aminoglycosides used in multiple courses or in combination with loop diuretics

Syndromes associated with progressive hearing loss:

- Neurofibromatosis, Osteopetrosis, Usher's Syndrome